



ISSN : 2320 4850

BI
MONTHLY

Asian Journal of Pharmaceutical Research And Development

(An International Peer Reviewed
Journal of Pharmaceutical
Research and Development)

A
J
P
R
D



Volume - 01

Issue - 05

SEP-OCT 2013

website: www.ajprd.com
editor@ajprd.com



Research Article

DESIGN AND DEVELOPMENT OF CHRONOPHARMACEUTICAL DRUG DELIVERY OF LANSOPRAZOLE

M. Sandeep^{*1}, V. Sai Kishore, B. Sudheer¹, S. Ershad, K. Adithya¹, D.S
Phanil kumar².

¹ Bapatla College of Pharmacy, Bapatla, Andhra Pradesh, India.

² Siddhartha institute of pharmacy, Jonnalagadda, Narsaraopeta, Andhra Pradesh, India.

Received: 8 Sept. 2013,

Revised and Accepted: 24 Oct. 2013

ABSTRACT

The aim of the present investigation is to develop a pulsatile drug delivery system based on an insoluble capsule body filled with Lansoprazole matrix tablet and sealed with HPMC K 100 plug. The Lansoprazole matrix tablets were prepared by wet granulation method. The plugs of varying thickness and hardness were prepared by direct compression which was then placed in the capsule opening. The drug delivery system was designed to deliver the drug at such a time when it was needed (nocturnal time). Dissolution studies of pulsatile capsule device in media with different pH (1.2, 7.4 and 6.8) showed that drug release in colon could be modulated by optimizing the concentration of polymers in the plug and also the position of plug. The study showed that, lag time prior to drug release was highly affected by the plug position. The dissolution data revealed that the plug position and the composition of plug were very important to achieve an optimum formulation. Drug-polymer interaction studies indicated no interaction or complexation in between the drug and the polymer.

KEYWORDS: Lansoprazole, wet granulation method, Hydrogel plug, Pulsing cap, Stability studies.

INTRODUCTION

A chronotherapeutic agent represents a pharmaceutical product that contains a dynamic element such as a delivery system to deliver the drug at the time when it is needed. A pulsatile drug delivery system is characterized by a lag time that is an interval of no drug release followed by rapid or controlled or sustained drug release. The interest in pulsatile delivery has been developing in close connection with emerging chronotherapeutic views.

In this respect, it is by now well established that the symptoms of many pathologies are subject to circadian variation patterns. Hence, the possibility of accomplishing effective drug levels in accordance with the specific temporal requirements of an illness state holds considerable appeal in that it could improve therapeutic outcome.

These systems are beneficial for drugs having high first pass effect; drugs administered for diseases that follow chronopharmacological behavior, drugs having specific absorption site in GIT, targeting to colon, to increase the stability of dosage form and cases where night time dosing is required.

Lansoprazole is an antiulcer drug belonging to the class of proton pump inhibitor. Lansoprazole is a benzimidazole sulfoxide derivative and produces long lasting inhibition

For correspondence:

M Sandeep

C/O: Bapatla college of pharmacy,

Bapatla, Guntur (district),

Andhra Pradesh, Pin No: 522101.

Cell No: 9014364835.

Email: sandeep2k8@gmail.com.

of gastric acid secretion. Lansoprazole is effective in the treatment of duodenal or gastric ulcer, gastro esophageal reflux disease and in the treatment of Zollinger-Ellison syndrome.

In Peptic ulcer patients, pain, gastric distress and acute exacerbation of the disease are most likely in the late evening and early morning hours. Ulcer pain typically occurs after stomach emptying, following daytime meals and in the very early morning, disrupting sleep. This is attributed to high gastric secretion and slows gastric motility and emptying at night. Suppression of nocturnal acid is an important factor in duodenal ulcer healing. Once daily nocturnal administration of proton pump inhibitor medications not only reduce acid secretion more effectively but also promotes ulcer healing and reduce ulcer occurrence.

The rationale of this study is to design and evaluate an oral site-specific, pulsatile drug delivery system containing Lansoprazole, which can be targeted to colon in a pH and time dependent manner, to modulate the drug level in synchrony with the circadian rhythm of nocturnal acid secretion. In the present research work, we have attempted to develop a novel dosage form by using a chronopharmaceutical approach.

A pulsatile 'Tablet in Capsule' dosage form, taken at bed time with a programmed start of drug release early in morning hours, can prevent a sharp increase in the incidence of high gastric acid secretion during the early morning hours, a time when the risk of peptic ulcer is the greatest.

In the present study, polymers such as Xanthan gum, Guar gum, are selected for the preparation of matrix tablets of Lansoprazole by wet granulation method and isopropyl alcohol as granulating agent [1, 2].

MATERIALS AND METHODS

MATERIALS

Lansoprazole was obtained as a gift sample from Hetero labs (Hyderabad), Guar gum and Xanthan gum was gifted by Yarrow Chem. Products (Mumbai). All other excipients used were of analytical grade.

PREPARATION OF CROSS-LINKED GELATIN CAPSULES:

The '0' sized hard gelatin capsules; about 100 in number were taken. The bodies of the capsules were then placed on a wire mesh. 25ml of 15% v/v formaldehyde was taken into a desiccators and a pinch of potassium permanganate was added to it to generate formalin vapours. The reaction was carried out for 12 hours. After which the bodies were removed and dried at 50°C for 30 minutes to ensure completion of reaction between gelatin and formaldehyde vapour. The bodies were dried at room temperature to facilitate removal of residual formaldehyde [3].

PREPARATION OF LANSOPRAZOLE GRANULES AND TABLETS:

Lansoprazole granules were prepared by using different drug: polymer ratios each respectively as per formulae given in Table I. The tablets were formulated by employing wet granulation method using PVP K-30 as binder, isopropyl alcohol as granulating fluid, magnesium stearate as lubricant and talc as glidant. The required quantities of Lansoprazole, polymer, lactose, and PVP K-30 were weighed and mixed well. Then it was made into damp mass using a mixture of isopropyl alcohol as granulating fluid. The resulting damp masses were screened by passing them manually through a 12 no. mesh size and dried for 30 minutes at 35±0.5°C in the oven and then screened through a 16 no. mesh and then dried to constant weight in the oven. The granules were then mixed with the required quantities of lubricants, glidant and then compressed into tablets using 5mm punches and dies on rotary tablet punching machine [4].

FLOW PROPERTIES OF GRANULES:

Table II contain evaluation values of granules. Granules were evaluated for Angle of repose, Bulk and Tapped densities, Percentage compressibility (Carr's index) and Hausner's ratio [4].

Table I: Composition of Lansoprazole matrix tablets with different ratios of natural polymers.

Formula	Xanthan gum			Guar gum		
	F ₁	F ₂	F ₃	F ₄	F ₅	F ₆
	1:1	1:2	1:3	1:1	1:2	1:3
Lansoprazole (mg)	30	30	30	30	30	30
Polymer(mg)	30	60	90	30	60	90
PVP K-30 (mg)	10	10	10	10	10	10
Lactose(mg)	75	45	15	75	45	15
Isopropyl alcohol (ml)	Q.S	Q.S	Q.S	Q.S	Q.S	Q.S
Magnesium stearate(mg)	3	3	3	3	3	3
Talc(mg)	2	2	2	2	2	2
Total(mg)	150	150	150	150	150	150

Table II: Evaluation tests of granules.

Formulation code	Angle of repose	Bulk density	Tapped density	Compressibility index	Hausssners ratio
F ₁	26.94±0.021	0.276±0.014	0.314±0.013	12.10±0.024	1.137±0.012
F ₂	25.6±0.031	0.350±0.012	0.408±0.011	14.21±0.022	1.161±0.014
F ₃	25.42±0.052	0.320±0.020	0.370±0.009	11.89±0.009	1.134±0.017
F ₄	26.85±0.024	0.319±0.005	0.362±0.021	11.87±0.017	1.130±0.024
F ₅	27.01±0.035	0.351±0.009	0.393±0.019	10.68±0.014	1.119±0.014
F ₆	25.76±0.05	0.255±0.025	0.291±0.005	12.37±0.024	1.142±0.014

EVALUATION OF TABLETS:

Table III contain evaluation values of tablets. Tablets were evaluated for Tablet thickness, Tablet weight variation, Friability, Hardness, and drug content [4].

Drug content:

The drug content in each formulation was determined by triturating 20 tablets and powder equivalent to average weight was added in 100 ml of 6.8 pH phosphate buffer, followed by stirring. The solution was filtered through a 0.45μ membrane filter, diluted suitably and the absorbance of resultant solution was measured spectrophotometrically at 285 nm using 6.8 pH phosphate buffer as blank.

PREPARATION AND EVALUATION OF HYDROGEL PLUG:

Four types of plugs were prepared by compressing polymer: lactose (1:1) ratio using 7 mm punches and dies on rotary tablet punching machine. The hydrogel plugs were evaluated for thickness, hardness, and lag time parameters. The composition and evaluation values of different types of plugs were given in the Table IV and Table V respectively. Fig. 1 represents the lag times of different hydrogel plugs [5].

DESIGNING OF PULSINCAP:

The Pulsincap was similar in appearance to a hard gelatin capsules, but the main body was water insoluble. Lansoprazole tablets were placed into the formaldehyde treated bodies by hand filling. The capsules containing the tablets were then plugged with optimized hydrogel plug.

The joint of the capsule body and cap was sealed with a small amount of the 5% ethyl

cellulose ethanolic solution. The sealed capsules were completely coated by dip coating method with 5% CAP in 8:2 (v/v) mixture of acetone: ethanol, plasticized with dibutylphthalate (0.75%), to prevent variable gastric emptying. Coating was repeated until an 8–12% increase in weight is obtained. % weight gain of the capsules before and after coating was determined [6].

Table III: Evaluation tests of tablets.

Parameters	Formulations					
	F ₁	F ₂	F ₃	F ₄	F ₅	F ₆
Tablet thickness(mm)	3.1±0.01	3.6±0.05	3.4±0.03	3.7±0.01	3.2±0.04	2.10±0.05
Hardness	4.7±0.021	4.5±0.025	4.8±0.032	4.6±0.038	5.1±0.042	5.0 ±0.01
Tablet weight variation(mg)	150±1	150±4	150±1	150±2	150±1	150±4
Friability	0.40±0.010	0.34±0.018	0.45±0.024	0.61±0.036	0.67±0.048	0.69 ±0.022
Drug content	100.14±0.13	99.78±0.15	99.56±0.11	100.1±0.38	100.78±0.87	99.67 ±0.08
Dissolution studies	99.24±0.06	99.68±0.04	99.31±0.09	99.60±0.03	99.32±0.09	99.04±0.08

Table IV: Composition of hydrogel plugs.

Hydrogel plugs	Ingredients (1:1)	Quantity (mg)
HP1	HPMC K-100:Lactose	100
HP2	Carbapol:Lactose	100
HP3	Na CMC:Lactose	100
HP4	Methyl Cellulose:Lactose	100

IN-VITRO DISSOLUTION PROFILE OF PULSATILE CAPSULE CONTAINING LANSOPRAZOLE MATRIX TABLET:

Dissolution studies were carried out by using USP XXIII dissolution test apparatus. Capsule was tied to paddle with a cotton thread so that the capsule should be immersed completely in dissolution media but not float. In order to simulate the pH changes along the GI tract, three dissolution media with pH 1.2, 7.4 and 6.8 were sequentially used referred to as sequential pH change method. When performing experiments, the pH 1.2 medium was first used for 2 hours (since the average gastric emptying time is 2 hrs.), then removed and the fresh pH 7.4 phosphate buffer saline was added. After 3 hours (average small

intestinal transit time is 3 hrs.), then the medium was removed and colonic fluid pH 6.8 buffer was added for subsequent hours. 900ml of the dissolution medium was used at each time. Rotation speed was 100 rpm and temperature was maintained at 37±0.5°C. 5 ml of dissolution media was withdrawn at predetermined time intervals and fresh dissolution media was replaced. The withdrawn samples were analyzed at 285 nm, by UV absorption spectroscopy and the cumulative percentage release was calculated over the sampling times [7].

Fig.2 and 3 shows the percentage drug release profile of pulsatile capsule containing Lansoprazole matrix tablets prepared with xanthan gum and guar gum.

DRUG-EXCIPIENT INTERACTION STUDIES:

Fourier Transform Infrared (FTIR) Spectroscopy studies were used for the evaluation of physicochemical compatibility and interactions, which helps in the prediction of interaction of the drug with Xanthan gum, Guar gum, diluents and lubricants used in tablet formulations. In the present study 1:1 ratio was used for preparation of physical mixtures and analyzed for compatibility

studies. Fig.4, 5, and 6 were the FTIR graphs of Lansoprazole, physical mixture of Lansoprazole and xanthan gum, and physical mixture of Lansoprazole and guar gum respectively [8].

STABILITY STUDIES:

Stability studies as per ICH guidelines were carried out for optimized formulation. The stability studies were carried out at $25 \pm 20^\circ\text{C}$ / $60 \pm 5\%$ RH and $40 \pm 20^\circ\text{C}$ / $75 \pm 5\%$ RH for 3 months [9].

Table V: Evaluation characteristics of hydrogel plugs.

Hydrogel Plug Code	Thickness(mm)	Hardness (kg/cm ²)	Lag time (hours)
HP1	3.16	4.5	5
HP2	3.29	4.2	4.5
HP3	3.24	3.8	4
HP4	3.54	3.5	3

RESULTS AND DISCUSSIONS

Pulsincap dosage form was a capsule which consists of a water insoluble body and a water soluble cap. The drug formulation (matrix tablet) was sealed within the capsule body by means of a hydrogel plug. The primary reaction of formaldehyde with gelatin (main constituent of capsule) probably is the formation of methylol amines. The tannin effect is due to a condensation reaction which transforms the methyl group into cross linking methylene bridges. The capsule bodies which were exposed to formaldehyde vapours for 12hrs were not dissolved in PH 7.4 phosphate buffer medium even after 48hrs (hardened capsule bodies were softened only after 24hrs). Thus for the present study, capsule bodies which were exposed to formaldehyde vapors to 12hrs were chosen for the preparation of pulsincaps. It was sealed with unhardened cap of the capsule.

The formulations fitted with the various hydrogel plugs HP1, HP2, HP3, HP4 shown 0.4%, 7.18%, 15.65% and 18.25% of drug release respectively at the end of 5th hour. It was observed that 100 mg hydrogel plug (HPMC K100 and lactose in 1:1 ratio) having

4.5 kg/cm² hardness was satisfactory to retard the drug release in small intestinal fluid and to eject out the plug in colonic fluid and releasing the matrix tablet into colonic fluid.

During dissolution studies, it was observed that, the enteric coat of the cellulose acetate phthalate was intact for 2 h in pH 1.2, but dissolved in intestinal pH, leaving the soluble cap of capsule, which also dissolved in pH 7.4, then the exposed polymer plug absorbed the surrounding fluid, swelled and released the drug through the swollen matrix. After complete wetting of the plug, it formed a soft mass, which was then easily ejected out of the capsule body; releasing the matrix tablet into simulated colonic fluid (pH 6.8 phosphate buffer). With all the formulations, there was absolutely no drug release in pH 1.2, thus indicating the efficiency of 5% CAP for enteric coating [9].

From the *In-vitro* release studies of device, it was observed that with all formulation, there was absolutely no drug release in simulated gastric fluid (acidic pH 1.2) for 2 hours and in simulated intestinal fluid (pH 7.4 phosphate buffer). Burst effect was found in colonic medium (pH 6.8 phosphate buffer). *In-vitro*

release profiles in colonic medium were found to have very good sustaining efficacy.

Pulsin caps loaded with matrix tablets prepared with Lansoprazole and xanthan gum in 1:1,1:2 and 1:3 ratios shown sustained drug release for a period of 8 hours (5th hour to 13th hour), 10 hours (5th hour to 15th hour) and 12 hours (5th hour to 17th hour) respectively.

Comparative *In-vitro* drug release profiles plot of Lansoprazole tablets prepared with xanthan gum in different ratios by wet granulation method were shown in Fig. 2.

Pulsin caps loaded with matrix tablets prepared with Lansoprazole and guar gum in 1:1,1:2 and 1:3 ratios shown sustained drug release for a period of 7.5 hours (5th hour to 12.5th hour), 9.5 hours (5th hour to 14.5th hour) and 11.5 hours (5th hour to 16.5th hour) respectively. Comparative *In-vitro* drug release profiles plot of Lansoprazole tablets prepared with xanthan gum in different ratios

by wet granulation method were shown in Fig. 3. The FTIR study reveals that the same characteristic features of pure drug were also observed in the drug loaded matrix tablets. Thus there was no significant drug polymer interaction which indicates the stable nature of the drug in all formulations.

The pulsincapsules were packed in the screw capped bottles and stored at $25 \pm 2^\circ\text{C}$, $60 \pm 5\%$ RH and at $40 \pm 2^\circ\text{C}$, $75 \pm 5\%$ RH for 3 months. Drug release from pulsincapsules before and after storage under varying conditions were evaluated periodically at the regular interval of every month. The results indicated that the drug release from the pulsincapsules was not changed significantly when stored at varying conditions. Thus the drug release from pulsincapsules was found to be quite stable.




Fig. 1: Plot of Lag times for hydrogel plugs.




Fig. 2: Comparative *In-vitro* drug release profile plot of Lansoprazole tablets prepared with xanthan gum in different ratios by wet granulation method.




Fig.3: Comparative *In-vitro* drug release profile plot of Lansoprazole tablets prepared with guar gum in different ratios by wet granulation method.




Fig.4: FTIR graph of Lansoprazole.




Fig.5: FTIR graph of formulation containing Lansoprazole and xanthan gum.




Fig.6: FTIR graph of formulation containing Lansoprazole and guar gum.

CONCLUSION

It is possible to release a drug over a predetermined period of time with specific release rates by controlling the polymers used to prepare plugs. The obtained results showed the capability of the system in delaying drug release for a programmable period of time and the possibility of exploiting such delay to attain colon targeting.

REFERENCES

1. Robert O. Williams III, David R. Taft, Jason T. McConville. *Advanced Drug Formulation Design to Optimize Therapeutic Outcomes*. New York: informa healthcare; 2008; 237-238.
2. Sungthangjeen S., Puttipipatkachorn S., Paeratakul O., Dashevsky A., Bodmeier R., Development of pulsatile release tablets with swelling and rupturable layers. *J Controlled Release*. 2004; 95(2):147-59.
3. Sukanya M, Sai Kishore V, Design and development of chronopharmaceutical drug delivery of simvastatin, *Journal of Chemical and Pharmaceutical Research*. 2012; 4(6): 3195-3200.
4. Sarojini Sarangapani, Manavalan Rajappan, Lansoprazole Release from a Floating Dosage Form based on the Natural Polymer of Delonix regia, *International Journal of PharmTech Research*. 2012; 4(3): 1084-1095.
5. Avinash R. Tekade, Surendra G, Gattin. Development and evaluation of pulsatile drug delivery system using novel polymers. *Pharmaceutical Development and Technology*. 2009; 14(4): 380-387.
6. Dasharath M Patel, Rushiraj H Jani, Chhagan N Patel. Design and evaluation of colon targeted modified pulsincap delivery of 5-fluorouracil according to circadian rhythm, *Int J Pharm Investig*. 2011; 1(3): 172-181.
7. Rongfeng Hu, Jiabi Zhu, Guang Chen, Yuliang Sun, Kangkang Mei, Shi Li. *AJPS*. 2006; 1: 47-52.
8. Nakhat P.D, Kondawar A.A, Rathil L.G, yeole P.G, Development and in vitro evaluation of mucoadhesive tablets of metoprolol tartrate, *Ind J Pharm Sci*. 2008; 70(1): 121-125.
9. Amit Bhat, Chowdary K.P.R, Shobharani R.H, *IJPSR*. 2011; 2(4): 1023-1030.