
Available online on 15.10.2026 at http://ajprd.com

Asian Journal of Pharmaceutical Research and Development

Open Access to Pharmaceutical and Medical Research

© 2013-25, publisher and licensee AJPRD, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Review Article

A Review: Potential Antihemorrhoidal Activity of Elephant Foot Yam (*Amorphophallus Paeoniifolius)*

Vaishnavi S. Shelke[1*], Jyoti R. Bhagat[2]

[1]Department of Pharmacy, Pravara Rural Education Society's College of Pharmacy (for Women), Chincholi, Nashik, 422101, (Maharashtra), India

[2]Department of Pharmaceutical Chemistry, Pravara Rural Education Society's College of Pharmacy (for Women), Chincholi, Nashik, 422101, (Maharashtra), India

ABSTRACT

Hemorrhoids, commonly known as piles, are a prevalent anorectal condition characterized by the enlargement and distal displacement of the normal anal cushions. Common symptoms include rectal bleeding, prolapse, itching, discomfort, and pain, particularly in patients with complicated disease. Dietary modification, adequate fiber intake, hydration, and appropriate medical or procedural interventions form the foundation of current management. Medicinal plants have also attracted interest because of their traditional applications and diverse phytochemical constituents. Elephant foot yam (*Amorphophallus paeoniifolius*), a tropical tuber belonging to the family Araceae, is widely consumed as a vegetable and has been used in traditional Indian medicine for various gastrointestinal and inflammatory conditions. Its corm contains carbohydrates, dietary fiber, minerals, and reported classes of secondary metabolites, including phenolic compounds, flavonoids, tannins, saponins, and sterol- or terpenoid-related compounds. These constituents provide a basis for investigating its possible biological activities, including antioxidant, anti-inflammatory, and astringent effects. However, the presence of these constituents does not establish clinical efficacy against hemorrhoids. Furthermore, calcium oxalate crystals in raw or inadequately processed corms can cause irritation. This review discusses hemorrhoidal disease, its anatomy, classification, etiology, pathophysiology, and management, followed by the botanical characteristics, chemical constituents, traditional uses, and potential antihemorrhoidal relevance of elephant foot yam. The review highlights the need for standardized phytochemical analysis, validated experimental models, safety assessment, and clinical studies before the plant can be recommended as a therapeutic agent for hemorrhoids.

Keywords: Hemorrhoids, Piles, *Amorphophallus paeoniifolius*, elephant foot yam, phytochemicals, flavonoids, tannins, anti-inflammatory activity, and medicinal plants.

ARTICLE INFO: Received 23 March 2026; Review Complete 18 July 2026; Accepted 14 Sept. 2026; Available online 15 Oct. 2026

Cite this article as:

Shelke VS, Bhagat JR, A Review: Potential Antihemorrhoidal Activity Of Elephant Foot Yam (*Amorphophallus Paeoniifolius)*, Asian Journal of Pharmaceutical Research and Development. 2026; 14(5): 159-167 DOI: **http://dx.doi.org/10.22270/ajprd.v14i5.1879**

*Address for Correspondence:

Vaishnavi S. Shelke, Department of Pharmacy, Pravara Rural Education Society's College of Pharmacy for Women), Chincholi, Nashik, 422101, (Maharashtra), India

INTRODUCTION

The term *hemorrhoid* is derived from the Greek words *haima*, meaning blood, and *rhoos*, meaning flow. Hemorrhoids refer to symptomatic enlargement and distal displacement of the normal vascular cushions of the anal canal. These cushions are normal anatomical structures that contribute to continence and provide support during defecation. Hemorrhoidal disease develops when these structures undergo pathological enlargement, displacement, inflammation, or thrombosis [1,2].

Hemorrhoids are among the most common anorectal disorders and may present with bright-red rectal bleeding, prolapse, itching, mucus discharge, discomfort, or pain. Their development is associated with several factors, including altered bowel habits, prolonged straining, increased intra-abdominal pressure, and deterioration of the connective tissues supporting the anal cushions. The clinical presentation varies according to the type and severity of the disease [1–3].

Conventional management includes dietary modification, adequate fluid intake, fiber supplementation, avoidance of prolonged straining, and appropriate medical or procedural treatment. Patients whose symptoms persist despite conservative management may require office-based procedures or surgery, depending on the grade and clinical characteristics of the hemorrhoids [3].

Medicinal plants have historically been investigated for their potential roles in managing inflammatory conditions, digestive disorders, and minor bleeding. Their biological activities are often associated with secondary metabolites

such as flavonoids, tannins, saponins, and terpenoids. However, the pharmacological effects of individual compounds vary, and traditional use alone cannot establish therapeutic efficacy [4].

Elephant foot yam, botanically identified as *Amorphophallus paeoniifolius* (Dennst.) Nicolson, is an important tropical tuber crop belonging to the family Araceae. It is cultivated and consumed in several Asian countries, including India. Its corm is valued for its nutritional content and traditional medicinal applications. The plant has been discussed in the context of gastrointestinal health and other traditional remedies, including those used for piles [5,6].

The reported nutritional and phytochemical characteristics of elephant foot yam provide a rationale for investigating its possible relevance to hemorrhoidal disease. Nevertheless, direct evidence demonstrating that the plant or its extracts reduce hemorrhoidal bleeding, prolapse, inflammation, or pain in humans remains insufficient. Therefore, this review evaluates the existing background literature and identifies the research required to determine its potential antihemorrhoidal activity.

REVIEW OBJECTIVES

The objectives of this review are:

1. To describe the anatomy, classification, etiology, and pathophysiology of hemorrhoidal disease.
2. To summarize conventional approaches to the management of hemorrhoids.
3. To describe the botanical characteristics, distribution, and traditional medicinal uses of elephant foot yam.
4. To review the major nutritional components and reported phytochemical constituents of *Amorphophallus paeoniifolius*.
5. To discuss the possible relevance of these constituents to anti-inflammatory, antioxidant, astringent, and bowel-regulating effects.
6. To identify safety concerns, limitations in the available evidence, and future research requirements.

REVIEW METHODOLOGY

This article is structured as a narrative review using the literature and references identified in the supplied material. The sources include review articles, clinical practice guidelines, phytochemical investigations, and publications concerning the morphology, cultivation, and medicinal applications of elephant foot yam.

The discussion integrates established information on hemorrhoidal disease with preliminary information on the plant's phytochemical profile. It is not a systematic review, and no formal database search, screening protocol, or quantitative synthesis has been performed for this draft. Consequently, the discussion should be interpreted as a narrative synthesis rather than a comprehensive evaluation of all available evidence.

HEMORRHOIDAL DISEASE


Definition

Hemorrhoids are vascular and connective-tissue cushions situated within the anal canal. In their normal state, they contribute to the maintenance of anal continence and protect the anal canal during defecation. Hemorrhoidal disease refers to symptoms resulting from enlargement, displacement, inflammation, or thrombosis of these structures [1,2].

Anatomy and physiological functions

The principal anal cushions are commonly described in the left lateral, right anterior, and right posterior positions of the anal canal. They contain vascular channels, connective tissue, smooth muscle, and supporting elastic fibres. The vascular structures include specialized arteriovenous communications and venous sinusoids supplied by branches of the rectal arterial system [1,2].

These cushions contribute to the resting closure of the anal canal and provide additional padding during continence. Their supporting connective tissues help maintain their anatomical position. Degeneration or weakening of these supporting structures, together with vascular changes, may contribute to their enlargement and downward displacement [1,2].



Types of hemorrhoids

Hemorrhoids are classified according to their relationship with the dentate line.

Internal hemorrhoids

Internal hemorrhoids arise from the vascular cushions above the dentate line and are covered by rectal mucosa. They

commonly present with painless, bright-red rectal bleeding or prolapse. Pain may occur when complications such as strangulation or thrombosis develop.

External hemorrhoids

External hemorrhoids occur below the dentate line and are covered by anoderm. Because this region has somatic sensory innervation, external hemorrhoids can be painful, particularly when thrombosis occurs. Acute thrombosis may produce a tender, swollen perianal lump.

Mixed hemorrhoids

Mixed hemorrhoids involve both internal and external components. Treatment depends on the extent of prolapse, symptoms, external disease, and associated complications [2,3].

Goligher classification of internal hemorrhoids

The Goligher classification grades internal hemorrhoids according to the extent of prolapse and whether the prolapsed tissue can be reduced.

Grade	Clinical description	Common presentation	General treatment approach
I	Enlarged hemorrhoids without prolapse beyond the anal canal	Painless bleeding	Dietary modification, fiber supplementation, and symptom-directed treatment
II	Prolapse during defecation or straining, followed by spontaneous reduction	Bleeding, irritation, or prolapse	Conservative management; office procedures if symptoms persist
III	Prolapse requiring manual reduction	Prolapse, bleeding, mucus discharge, and discomfort	Office procedures or surgery, depending on severity and response
IV	Permanently prolapsed and irreducible hemorrhoids	Persistent prolapse, discomfort, and possible strangulation or thrombosis	Surgical assessment; excisional treatment is often considered

The classification assists clinicians in selecting appropriate treatment. However, treatment decisions should also consider symptom severity, external components, patient preferences, comorbidities, and previous treatment outcomes [2,3].

Etiology and risk factors

The precise cause of hemorrhoidal disease is multifactorial and is not completely understood. Important contributing factors include:

- **Altered bowel habits:** Constipation, hard stools, and repeated straining may contribute to displacement of the anal cushions. Diarrhea and frequent bowel movements may also aggravate anorectal symptoms.
- **Low dietary fiber intake:** Insufficient fiber can contribute to hard stools and difficult defecation.
- **Increased intra-abdominal pressure:** Pregnancy, prolonged straining, obesity, and certain abdominal conditions may increase pressure on the anorectal vascular structures.
- **Age-related tissue changes:** Deterioration of connective tissue may reduce the stability of the anal cushions.
- **Prolonged sitting and inactivity:** These may be associated with unfavorable bowel habits and other contributing factors.
- **Pregnancy and childbirth:** Hormonal changes, venous compression, constipation, and increased abdominal pressure during delivery may aggravate hemorrhoidal symptoms.
- **Other associated conditions:** Pelvic floor dysfunction and certain gastrointestinal or abdominal disorders may contribute to symptoms in selected patients.

These factors should be regarded as potential contributors rather than universal causes. The relationship between constipation and hemorrhoids is complex, and constipation alone does not explain all cases [2,4,5].

Pathophysiology

Hemorrhoidal disease involves structural and vascular changes in the normal anal cushions. Three interrelated mechanisms are particularly relevant.

Degeneration of connective tissue and downward displacement

The anal cushions are supported by connective tissue, elastic fibers, and smooth muscle. Age-related degeneration and repeated mechanical stress may weaken this supporting framework. As the supporting tissues lose their integrity, the cushions may move distally and protrude during defecation [5].

Vascular congestion and altered blood flow

Changes in vascular filling, arterial inflow, venous drainage, and local hemodynamics may contribute to enlargement of the hemorrhoidal cushions. The specialized vascular architecture of the anorectal region helps explain why internal hemorrhoids may produce bright-red bleeding [1,5].

Inflammation and tissue remodeling

Repeated mechanical stress, mucosal irritation, and tissue injury may contribute to local inflammation and changes in extracellular matrix remodeling. These changes may worsen bleeding, swelling, and prolapse. Matrix metalloproteinases and other mediators have been discussed in relation to connective-tissue remodeling in hemorrhoidal disease [5,7].

Together, these mechanisms help explain the clinical features of hemorrhoids and provide a basis for investigating interventions that may influence inflammation, tissue irritation, or bowel habits. They do not, however, establish that a particular plant extract will be effective.

CONVENTIONAL MANAGEMENT OF HEMORRHOIDS

Dietary and lifestyle modification

Conservative management is the first-line approach for many patients with symptomatic hemorrhoids. It includes

increasing dietary fiber, maintaining adequate hydration, avoiding excessive straining, responding promptly to the urge to defecate, and reducing prolonged sitting on the toilet. Fiber supplementation can improve symptoms, particularly bleeding and overall discomfort [3,8].

Fiber-rich foods include vegetables, fruits, whole grains, and legumes. The quantity of fiber should be increased gradually and accompanied by adequate fluid intake when medically appropriate.

Symptomatic medical treatment

Warm sitz baths may provide temporary relief from discomfort. Selected topical preparations containing protectants, local anesthetics, astringents, or short-term corticosteroids may help relieve itching and irritation. Their benefits vary, and prolonged use of some topical products may cause adverse effects [8].

Certain oral phlebotonic preparations have been investigated for hemorrhoidal symptoms. Although some studies suggest improvement in bleeding or other symptoms, the strength and consistency of the evidence vary. Such preparations should not replace appropriate assessment or definitive treatment when indicated [3,8].

Office-based procedures and surgery

For internal hemorrhoids that remain symptomatic despite conservative management, office-based procedures such as rubber band ligation, injection sclerotherapy, or infrared coagulation may be considered. More advanced prolapse, significant external disease, or persistent symptoms may require surgical evaluation. Excisional hemorrhoidectomy is an established option for selected patients with advanced or combined disease [3].

Emerging minimally invasive procedures

Laser hemorrhoidoplasty and Doppler-guided hemorrhoidal artery ligation are among the procedures investigated for selected patients. The choice of procedure depends on disease characteristics, available expertise, expected benefits, risks, and the quality of supporting evidence. These approaches should not be presented as universally superior to conventional treatments [3].

ELEPHANT FOOT YAM (*AMORPHOPHALLUS PAEONIIFOLIUS*)

Botanical description

Elephant foot yam, *Amorphophallus paeoniifolius* (Dennst.) Nicolson, belongs to the family Araceae. It is a tropical tuber crop cultivated primarily for its edible underground corm. The plant is valued for its agricultural productivity, nutritional contribution, and traditional uses [6,9].

The corm is rich in starch and supplies energy. It also contains dietary fiber and smaller quantities of proteins and minerals. Its composition varies with cultivar, growing conditions, maturity, storage, and processing methods [6,9].



Taxonomic classification

Taxonomic rank	Classification
Kingdom	Plantae
Division	Tracheophyta
Class	Liliopsida
Order	Alismatales
Family	Araceae
Genus	*Amorphophallus*
Species	*Amorphophallus paeoniifolius*

The table provides a practical botanical classification. Exact intermediate taxonomic ranks may differ among taxonomic databases.

Distribution

Amorphophallus paeoniifolius is associated with tropical and subtropical regions of Asia and the Pacific. It is cultivated in

India and other Asian countries, where its corm is consumed as a vegetable. In India, it is grown in several states, including Kerala, Tamil Nadu, Andhra Pradesh, Maharashtra, Odisha, West Bengal, and parts of eastern and northern India [6,9].

Traditional medicinal uses

Elephant foot yam has a history of use in traditional Indian medical systems, including Ayurveda, Siddha, and Unani. The corm has been used traditionally for certain digestive complaints and other conditions, and traditional literature describes its use in remedies for piles [6,9].


These applications are relevant to ethnopharmacological research because they may help identify potential therapeutic uses for further investigation. However, traditional claims should not be interpreted as proof of clinical efficacy. Reliable evidence from standardized extracts, experimental studies, and controlled clinical trials is needed before recommending elephant foot yam as a treatment for hemorrhoids.

Nutritional significance

The nutritional profile of elephant foot yam is relevant to its use as a food crop.

- **Carbohydrates:** Starch is a major component and provides dietary energy.
- **Dietary fiber:** Contributes to total fiber intake and may support normal bowel function as part of a balanced diet.
- **Proteins:** Present in smaller amounts than carbohydrates.
- **Minerals:** The corm may provide minerals such as calcium, phosphorus, and iron, with quantities depending on the sample and analytical method.
- **Other nutrients:** Composition varies according to cultivation, maturity, and processing.

Dietary fiber has an established role in conservative hemorrhoid management. However, the contribution of elephant foot yam to daily fiber intake does not demonstrate a plant-specific antihemorrhoidal effect. Its nutritional value should be distinguished from the pharmacological activity of concentrated extracts [3,6,9].



CHEMICAL CONSTITUENTS OF ELEPHANT FOOT YAM

Phytochemical investigations of *Amorphophallus paeoniifolius* have reported several classes of primary and secondary metabolites. The identity and quantity of these constituents vary with the plant material, extraction solvent, analytical technique, geographical source, and processing conditions. Therefore, not every reported constituent will necessarily be present in every corm sample or extract [9,10].

Primary metabolites

- **Carbohydrates and starch**

  Carbohydrates, particularly starch, form a major part of the corm. These compounds primarily contribute to its nutritional and energy value.

- **Dietary fiber and polysaccharides**

  The corm contains dietary fiber and other polysaccharide components. Fiber is relevant to hemorrhoidal management because increased dietary fiber can improve stool consistency and reduce straining in some patients. The extent to which specific polysaccharides in elephant foot yam contribute to these effects requires appropriate compositional and physiological studies [3,9].

- **Proteins and amino acids**

  Proteins and amino acids contribute to the nutritional composition of the corm. Their concentrations may vary according to cultivar and growing conditions [9].

- **Lipids and fatty acids**

  Lipids occur in smaller quantities than starch. The types and concentrations of fatty acids should be established using appropriate analytical methods before assigning them specific pharmacological effects.

Secondary metabolites

- **Flavonoids and other phenolic compounds**

  Flavonoids and phenolic compounds are important phytochemical groups investigated for antioxidant and other biological activities. Some plant-derived flavonoids

can influence oxidative processes and inflammatory signaling in experimental systems.

Their relevance to hemorrhoidal disease is indirect: antioxidant or anti-inflammatory activity in a laboratory does not establish that a plant extract will reduce hemorrhoidal bleeding, prolapse, or pain. Individual compounds and their concentrations in *A. paeoniifolius* require analytical confirmation [9,10].

- **Tannins**

Tannins are polyphenolic compounds associated with astringent activity. They can interact with proteins and may produce a localized astringent effect under appropriate conditions. Because astringents are used in some topical preparations for symptom relief, tannins are of potential research interest.

Nevertheless, the presence of tannins in an extract does not establish that the extract is safe for anorectal application or capable of stopping hemorrhoidal bleeding. These effects must be investigated using appropriate pharmacological and safety studies [4,10].

- **Saponins**

Saponins are a diverse group of glycosides reported in various medicinal plants. Some saponins demonstrate anti-inflammatory or other biological activities in experimental models. Their occurrence and abundance in a particular elephant foot yam preparation must be verified before any therapeutic inference is made [9,10].

- **Alkaloids**

Alkaloids are nitrogen-containing compounds with diverse biological activities. Preliminary phytochemical investigations may identify alkaloid-like constituents in plant extracts; however, qualitative screening alone does not establish the identity, concentration, safety, or pharmacological significance of individual alkaloids.

- **Sterols and terpenoid-related compounds**

Plant sterols and terpenoid-related compounds are of interest in phytochemical research because some members of these chemical classes have demonstrated biological activities in other plant systems. Their presence, exact identity, and concentration in *A. paeoniifolius* should be confirmed using suitable chromatographic and spectrometric methods before specific pharmacological claims are made [9,10].

- **Glycosides**

Glycosides comprise a broad group of compounds in which a sugar is linked to another chemical group. The biological effects depend on the structure of the individual compound. The general detection of glycosides does not justify attributing a specific therapeutic action to an elephant foot yam preparation.

- **Calcium oxalate crystals and acridity**

Calcium oxalate crystals, including needle-shaped crystals known as raphides, are important antinutritional and irritant components of raw or inadequately processed elephant foot yam. They can cause itching, burning, and irritation of the oral mucosa and throat.

Appropriate culinary processing, including thorough cooking, is important to reduce acridity and improve edibility. Processing methods should be evaluated for their effectiveness in reducing oxalate content, maintaining nutritional quality, and preserving any phytochemicals of interest [9].

It is not scientifically justified to describe calcium oxalate crystals themselves as established anti-inflammatory agents in this context. Nor should enzyme treatment or fermentation be considered validated pharmaceutical processing methods without experimental evidence demonstrating oxalate reduction, phytochemical retention, microbiological safety, and reproducibility.

Summary of chemical constituents and their possible relevance

Chemical constituent or group	General role or reported property	Possible relevance to hemorrhoid research	Important limitation
Starch and carbohydrates	Energy source	Primarily nutritional	Not a direct antihemorrhoidal mechanism
Dietary fiber and polysaccharides	Support dietary fiber intake and bowel function	May be relevant to stool consistency when consumed as part of a suitable diet	Plant-specific therapeutic effects are unproven
Flavonoids and phenolics	Antioxidant activity in various experimental systems	Potential research interest in inflammatory processes	Requires compound identification and activity testing
Tannins	Astringent properties in some preparations	Potential relevance to topical symptom-relief research	Safety and hemostatic effects are unestablished for this plant
Saponins	Diverse biological activities reported across plant species	Potential pharmacological research interest	Composition and efficacy must be confirmed
Alkaloids	Structurally diverse secondary metabolites	Possible leads for compound-specific research	Identity, dose, and safety require confirmation
Sterols and terpenoids	Diverse biological activities among individual compounds	Potential leads for pharmacological screening	Specific compounds should not be assumed without analytical evidence
Glycosides	Diverse structural and biological group	May warrant further phytochemical investigation	No single therapeutic effect can be inferred for the whole class
Calcium oxalate crystals	Irritant and antinutritional factors in inadequately processed corms	Important for safety and processing evaluation	Not an established therapeutic constituent

The constituents listed above should be interpreted as reported or candidate phytochemical groups, not as a confirmed quantitative profile of every elephant foot yam preparation. Compound-specific identification should be supported by validated analytical methods.

POTENTIAL ANTIHEMORRHOIDAL ACTIVITY OF ELEPHANT FOOT YAM

Rationale for investigation

The traditional use of elephant foot yam for piles, together with the broader biological properties associated with some plant-derived compounds, provides a rationale for investigating its potential relevance to hemorrhoidal disease. However, the evidence must be considered at three distinct levels: traditional use, experimental pharmacological plausibility, and demonstrated clinical efficacy.

Traditional use can guide research, while laboratory studies can identify possible mechanisms. Neither level alone establishes that the plant is effective in patients.

Potential role of dietary fiber

Dietary fiber is one of the most directly relevant aspects of elephant foot yam's nutritional profile. Adequate fiber intake can improve stool consistency and reduce the need for excessive straining in patients with hemorrhoidal symptoms. Fiber supplementation is recommended as part of conservative management [3,8].

Elephant foot yam may contribute to dietary fiber intake when properly cooked and consumed as part of a balanced diet. Nevertheless, the total fiber content of a particular serving, the individual's overall dietary pattern, and tolerance must be considered. There is insufficient evidence to conclude that eating elephant foot yam specifically treats hemorrhoids or that it is superior to other fiber-containing foods.

Potential anti-inflammatory activity

Inflammatory processes may contribute to local irritation and tissue changes in hemorrhoidal disease. Certain flavonoids, phenolic compounds, and saponins from medicinal plants have been investigated for their potential effects on inflammatory pathways [4].

If validated extracts of elephant foot yam demonstrate relevant anti-inflammatory activity in appropriate experimental models, this could justify further investigation. Such activity should not be presumed solely from the presence of a phytochemical class. The active compounds, effective concentrations, toxicity, and relevance to anorectal tissues would need to be established.

Potential astringent and hemostatic relevance

Tannins from certain plants can produce astringent effects through interactions with proteins. This property provides a theoretical basis for investigating selected plant extracts in topical preparations intended for minor irritation.

However, hemorrhoidal bleeding may arise from vascular and mucosal changes that require proper diagnosis. No adequate evidence presented in the sources reviewed here establishes that elephant foot yam or its extracts reliably stop hemorrhoidal bleeding. Astringency must not be equated with proven hemostatic efficacy.

Potential antioxidant activity

Oxidative stress is investigated in many inflammatory and tissue-remodeling conditions. Some phenolic compounds and flavonoids demonstrate antioxidant activity in laboratory assays. Such findings may provide preliminary information about a plant's biological properties.

Antioxidant activity measured in a chemical assay, however, does not demonstrate absorption, activity at the affected tissue, or clinical benefit. Further studies should determine whether standardized elephant foot yam extracts exhibit reproducible antioxidant effects at non-toxic concentrations and whether these effects have any meaningful relationship to hemorrhoidal disease.

Overall assessment of the evidence

The potential antihemorrhoidal relevance of elephant foot yam remains exploratory. Its traditional use and nutritional contribution justify further research, but the evidence described in this review does not establish it as an effective treatment for hemorrhoids.

In particular, the available information does not establish an effective dose, a validated medicinal formulation, a reliable route of administration, or comparative clinical benefit. The plant should therefore not replace guideline-based conservative care, office procedures, or surgery when these are clinically indicated.

SAFETY AND PHARMACEUTICAL DEVELOPMENT CONSIDERATIONS

Plant material identification and standardization

Research should begin with authenticated plant material, documented collection details, and a verified botanical identity. The corm should be processed under standardized conditions, and the final preparation should be characterized for its physicochemical properties and phytochemical composition.

Reduction of oxalate-related irritation

Raw or inadequately processed elephant foot yam may cause significant irritation because of calcium oxalate crystals. Any proposed medicinal preparation must undergo validated processing and testing to determine residual oxalate content and safety. Traditional culinary preparation should not automatically be considered suitable for pharmaceutical manufacturing.

Phytochemical retention and analytical validation

Potential processing methods include controlled soaking, cooking, and other appropriately validated procedures. The effects of each method on oxalate content and target phytochemicals should be measured rather than assumed.

For experimental pharmaceutical development, suitable analyses may include:

- Preliminary phytochemical screening.
- Total phenolic and total flavonoid assays.
- Chromatographic profiling using HPLC or LC-MS, where appropriate.
- Validated oxalate estimation.
- Assessment of extract stability and batch-to-batch consistency.
- Appropriate microbiological and contaminant testing.

Enzymatic processing or fermentation may be investigated as experimental approaches, but their suitability cannot be established without validation of the relevant process parameters, safety, reproducibility, and final-product quality.

Toxicity and route of administration

A preparation intended for oral consumption may have a different safety profile from a concentrated extract intended for topical or anorectal application. Each proposed route requires appropriate safety evaluation.

Before human studies, researchers should evaluate relevant cytotoxicity, local irritation, acute and repeated-dose toxicity, and other risks according to the intended formulation and applicable research standards. Traditional dietary use does not establish the safety of concentrated extracts or direct application to anorectal tissues.

Requirements for clinical evaluation

If preliminary laboratory and safety studies demonstrate a sufficient rationale, subsequent research could investigate standardized preparations in appropriately designed clinical trials. Such studies should use ethically approved protocols, informed consent, predefined outcomes, and appropriate control groups.

Relevant outcomes may include the frequency and severity of bleeding, prolapse, pain, itching, adverse events, and the need for additional treatment. Disease severity should be documented, and participants should receive appropriate standard care. Until such evidence becomes available, efficacy claims should remain provisional.

RESEARCH GAPS AND FUTURE PROSPECTS

Several areas require further investigation before elephant foot yam can be considered a potential therapeutic candidate for hemorrhoidal disease.

1. **Phytochemical characterization:** Identification and quantification of the compounds actually present in authenticated corm material and standardized extracts.
2. **Mechanistic studies:** Evaluation of antioxidant, anti-inflammatory, astringent, and other relevant activities using validated experimental methods.
3. **Oxalate reduction:** Development and validation of processing methods that reduce irritant oxalate content while preserving selected compounds of interest.
4. **Safety assessment:** Investigation of toxicity, local irritation, interactions, and formulation-specific risks.
5. **Formulation development:** Establishment of a reproducible formulation, appropriate route of administration, stability, and quality-control parameters.
6. **Preclinical efficacy:** Evaluation in scientifically appropriate models, with clear endpoints and adequate controls.
7. **Clinical trials:** Carefully designed studies to determine whether a standardized preparation provides meaningful benefits beyond established conservative management.

A key research priority is to separate the established benefits of dietary fiber from any proposed pharmacological effects of concentrated elephant foot yam extracts. This distinction would help prevent unsupported claims and guide the design of more informative studies.

CONCLUSION

Hemorrhoids are a common anorectal condition associated with changes in the vascular cushions, weakening of their supporting connective tissues, and symptoms such as bleeding, prolapse, itching, and discomfort. Current management emphasizes dietary fiber, adequate hydration, avoidance of straining, symptom-directed treatment, and appropriate procedural or surgical intervention when required.

Elephant foot yam (*Amorphophallus paeoniifolius*) is a nutritionally important tropical tuber with a history of traditional medicinal use. Its corm contains starch, dietary fiber, proteins, minerals, and reported classes of phytochemicals, including flavonoids, phenolic compounds, tannins, saponins, and other secondary metabolites. These constituents provide a scientific rationale for further investigation of its nutritional and potential pharmacological properties.

Among its possible areas of relevance, dietary fiber has an established role in the conservative management of hemorrhoidal symptoms. In contrast, proposed anti-inflammatory, antioxidant, astringent, and hemostatic effects of elephant foot yam remain insufficiently established for the treatment of hemorrhoids. The presence of a particular phytochemical does not, by itself, demonstrate therapeutic effectiveness. In addition, calcium oxalate crystals create important processing and safety considerations.

In conclusion, elephant foot yam may be considered a candidate for further phytochemical and pharmacological investigation, rather than an established antihemorrhoidal medicine. Authenticated plant material, standardized extraction, validated analytical methods, toxicity testing, and well-designed clinical studies are necessary to determine whether it offers a safe and clinically meaningful benefit. Until such evidence is available, it should not replace established hemorrhoid management.

REFERENCES

1. Aigner F, Gruber H, Conrad F, Eder J, Wedel T, Zelger B, et al. Revised morphology and hemodynamics of the anorectal vascular plexus: impact on the course of hemorrhoidal disease. International Journal of Colorectal Disease. 2009;24(1):105–113.
2. Margetis N. Pathophysiology of internal hemorrhoids. Annals of Gastroenterology. 2019;32(3):264–272.
3. Hawkins AT, Davis BR, Bhama AR, Fang SH, Dawes AJ, Feingold DL, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Hemorrhoids. Diseases of the Colon & Rectum. 2024;67(5):614–623.
4. Rahimi R, Abdollahi M. Evidence-based review of medicinal plants used for the treatment of hemorrhoids. International Journal of Pharmacology. 2013;9(1):1–11. Available from: Article PDF.
5. Sandler RS, Peery AF. Rethinking what we know about hemorrhoids. Clinical Gastroenterology and Hepatology. 2019;17(1):8–15. Available from: Article PDF.
6. Singh A, Wadhwa N. A review on multiple potential of aroid: Amorphophallus paeoniifolius. International Journal of Pharmaceutical Sciences Review and Research. 2014;24(1):55–60.
7. Lohsiriwat V. Hemorrhoids: from basic pathophysiology to clinical management. World Journal of Gastroenterology. 2012;18(17):2009–2017.

8. Mott T, Latimer K, Edwards C. Hemorrhoids: diagnosis and treatment options. American Family Physician. 2018;97(3):172–179.

9. Suja G, Sundaresan S, John KS, Sreekumar J, Misra RS. Higher yield, profit and soil quality from organic farming of elephant foot yam. Agronomy for Sustainable Development. 2012;32(3):755–764.

10. De S, Dey YN, Ghosh AK. Phytochemical investigation and chromatographic evaluation of the different extracts of tuber of Amorphophallus paeoniifolius (Araceae). International Journal of Pharmaceutical and Biological Research. 2010;1(5):150–157.

11. Mahboubi M. Effectiveness of Myrtuscommunis in the treatment of hemorrhoids. Journal of Integrative Medicine. 2017;15(5):351–358.

12. Lorenzo-Rivero S. Hemorrhoids: diagnosis and current management. The American Surgeon. 2009;75(8):635–642.

13. Amorphophallus paeoniifolius (Dennst.) Nicolson. Botanical and traditional-use information. Additional botanical details should be verified against a recognized botanical database before submission.

14. Mott T, Latimer K, Edwards C. Hemorrhoids: diagnosis and treatment options. American Family Physician. 2018;97(3):172–179.